

bcl-6 Ab-2 (Clone BL6.02; same as PG-B6p)

Mouse Monoclonal Antibody

Cat. #MS-1114-S0, -S1, or -S (0.1ml, 0.5ml, or 1.0ml Supernatant)

Cat. #MS-1114-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)

Cat. #MS-1114-PCS (5 Slides) (Positive Control for Histology)

Description: Bcl-6 proto-oncogene product (a Kruppel-type zinc-finger protein) is mainly expressed in normal germinal center B cells and related lymphomas. Bcl-6 is involved in chromosome rearrangements at 3q27 in non-Hodgkin's lymphomas and Bcl-6 rearrangements have been detected in 33%-45% of diffuse large B cell lymphomas. Bcl-6 has been detected immunohistochemically in follicular lymphomas, diffuse large B cell lymphomas, Burkitt's lymphomas and in nodular, lymphocyte predominant Hodgkin's disease.

Epitope: aa 3-484

Species Reactivity: Human, Cow, Rabbit, Rat, Sheep, and Pig. Others-not tested

Clone Designation: BL6.02; same as PG-B6p

Ig Isotype / Light Chain: IgG₁ / κ

Immunogen:

Recombinant protein corresponding to amino acids 3-484 of the human Bcl-6 protein

Applications and Suggested Dilutions:

- Immunohistology (Formalin/paraffin)
(Use Ab at 1:20 for 60 min at RT)

*(Staining of formalin-fixed tissues REQUIRES boiling tissue sections in 1mM EDTA, pH 8.0 (**NEOMARKERS**® Cat. #AP-9004), for 10-20 min followed by cooling at RT for 20 min. and REQUIRES use of a high-sensitivity detection system such as UltraVision LP (Cat# TL-015))

The optimal dilution and incubation time for a specific application should be determined by the investigator.

Positive Control: Tonsil

Cellular Localization: Nuclear

Supplied As:

Tissue culture supernatant with 0.09% sodium azide,
or

Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

Storage and Stability:

Store vial at 4°C. When stored at 2-8°C, this antibody is stable for 24 months.

Key References:

1. Flenghi L, et al. (1996) Am J Pathol 148: 1543-55

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. NeoMarkers is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:


This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only

Thermo Fisher Scientific
Anatomical Pathology
47777 Warm Springs Blvd.
Fremont, CA 94539, USA
Tel: 1-510-991-2800
Fax: 1-510-991-2826
<http://www.thermo.com/labvision>



Manufactured by:
NeoMarkers
For
Lab Vision Corporation



Thermo Fisher Scientific
Anatomical Pathology
93-96 Chadwick Road, Astmoor
Runcorn, Cheshire WA7 1PR, UK
Tel: 44-1928-562600
Fax: 44-1928-562627
Labvision.uk@thermofisher.com