

Estrogen Receptor beta

Rabbit Polyclonal Antibody

Cat. #RB-10658-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)

Cat. #RB-10658-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)

Cat. #RB-10658-PCS (5 Slides) (Positive Control for Histology)

Description: ER exists in two types: alpha and Beta. They are similar in structure. The ligand binding domain and the DNA binding domain are highly conserved in these proteins while the N-terminal transactivating domain is diverged considerably. Five isoforms of the hER beta gene, designated hER beta 1-5 have been identified. ER-Beta binds to estrogen and activates genes through direct interaction with estrogen specific gene elements (ERE's). ER-Beta RNA has been detected in human thymus, spleen, ovary and testis and in rat ovary and prostate. This tissue distribution overlaps but is not identical to that of ER - alphaRNA.

Mol. Wt. Of Antigen: 53kDa

Epitope: C-terminal

Species Reactivity: Human. Others not tested.

Immunogen:

Synthetic peptide corresponding to the C-terminus of human ER beta.

Applications and Suggested Dilutions:

- Immunohistology (Formalin/paraffin)

(Ab at 1:25 for 30 min with LV's UltraVision)

* [Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.]

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: Breast Carcinoma

Cellular Localization: Nuclear

Supplied As: Affinity purified antibody fraction from rabbit anti-serum. Prepared in 10mM PBS, pH7.6, with 0.2% BSA and 15mM sodium azide. Or Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.


Storage and Stability: Store vial at 4°C. When stored at 2-8°C, this antibody is stable for 24 months.

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. NeoMarkers is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.



FFPE human breast carcinoma stained with anti-ERβ (cat.# RB-10658-P) using peroxidase conjugate and AEC chromogen. Note Nuclear staining.

For Research Use Only

