

D4-GDI

Rabbit Polyclonal Antibody

Cat. #RB-9042-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)

Cat. #RB-9042-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)

Cat. #RB-9042-PCS (5 Slides) (Positive Control for Histology)

Description: D4-GDI belongs to the family of Rho-GDIs or GDP-dissociation inhibitors for the Rho/Rac family of small G proteins; consisting of RhoGDI-1, D4-GDI/RhoGDI-2 and RhoGDI-3. It inhibits the release of GDP from the Rho proteins, essential for GTP-loading and activation of GTPase activity. D4-GDI is located in the cytoplasm and is involved in the regulation of the membrane association and dissociation cycle of Rho/Rac proteins. It is known to be a substrate for caspase 3 during Fas-mediated apoptosis.

Comments: This antibody is excellent for staining of formalin-fixed, paraffin-embedded tissues.

Mol. Wt. of Antigen: 28kDa

Epitope: Not determined

Species Reactivity: Human. Others not tested.

Immunogen:

Recombinant full-length human D4-GDI protein.

Applications and Suggested Dilutions:

- Western Blotting (Ab 1µg/ml for 2 hours at RT)
- Immunohistology (Formalin/paraffin)

(Ab 1:100 for 20 min with LV's LP and 30min with UV ONE or UV)

- [Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.]

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: Placenta and tonsil

Cellular Localization: Placenta and tonsil

Storage and Stability: Store vial at 4°C. When stored at 2-8°C, this antibody is stable for 24 months.

Supplied As: Affinity purified antibody fraction from rabbit anti-serum. Prepared in 10mM PBS, pH 7.6, with 0.2% BSA and 15mM sodium azide.

or

Pre-diluted antibody which is ready-to-use for immunohistochemical staining.


Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. NeoMarkers is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only



Human placenta stained with Anti-D4-GDI, Cat# RB-9042-P

